

Counterexample to the passive topological censorship of $K(\pi, 1)$ prime factors

Gregory A. Burnett

Department of Physics, University of Florida, Gainesville, Florida 32611
(5 April 1995)

A globally hyperbolic asymptotically flat spacetime is presented (having non-negative energy density and pressures) that shows that not all $K(\pi, 1)$ prime factors of the Cauchy surface topology are passively censored according to asymptotic observers in contradiction to an argument of Friedman, Schleich, and Witt.

04.20.Gz, 04.20.Cv

I. INTRODUCTION

One of the intriguing features of Einstein's theory of gravity is that it allows for the possibility that our Universe is topologically non-Euclidean. For instance, somewhere a wormhole may link otherwise very distant regions of space together. Why then have such structures not been observed? One possibility is that our Universe just happens to be topologically trivial. However, Friedman, Schleich, and Witt argue that even if the Universe is topologically nontrivial, observers that remain in the asymptotic region (i.e., do not venture into black holes) cannot probe this topology [1,2]. They name this effect *topological censorship*.

Before discussing their results, we first present a few definitions. Fix an asymptotically flat spacetime (M, g_{ab}) and denote the null infinity associated with an asymptotic region of this spacetime by $\mathcal{J}$ (which is the disjoint union of a past part $\mathcal{J}^-$ and a future part $\mathcal{J}^+$) [4]. Denote the universal covering spacetime associated with (M, g_{ab}) by $(\tilde{M}, \tilde{g}_{ab})$. If M is not simply connected, then $(\tilde{M}, \tilde{g}_{ab})$ will have many distinct null infinities $\mathcal{J}_i$ associated with $\mathcal{J}$.

An asymptotically flat spacetime (M, g_{ab}) is said to satisfy *active topological censorship* if there are no homotopically distinct causal curves from $\mathcal{J}^-$ to $\mathcal{J}^+$ [1]. In such a spacetime, there is no way an observer in the asymptotic region can actively probe the nontrivial topology of the spacetime by sending and receiving causal signals as the paths taken by any two such signals are necessarily homotopic. If M is simply connected, then active topological censorship is automatically satisfied. Otherwise, active topological censorship is equivalent to the condition that

$$\mathcal{J}^+(\mathcal{J}_i^-) \cap \mathcal{J}^-(\mathcal{J}_j^+) = \emptyset, \text{ for } i \neq j. \quad (1.1)$$

An asymptotically flat spacetime (M, g_{ab}) is said to satisfy *passive topological censorship* if there are no points $p \in M$ for which there exist homotopically distinct causal curves from p to $\mathcal{J}^+$ [1]. In such a spacetime, there is no way an observer in the asymptotic region can passively observe the nontrivial topology of the spacetime by receiving causal signals from some point in the spacetime

as the paths taken by any two such signals are necessarily homotopic. Again, if M is simply connected this is automatically satisfied. Otherwise, passive topological censorship is equivalent to the condition that

$$\mathcal{J}^-(\mathcal{J}_i^+) \cap \mathcal{J}^-(\mathcal{J}_j^+) = \emptyset, \text{ for } i \neq j. \quad (1.2)$$

Friedman, Schleich, and Witt have proven that active topological censorship holds for the globally hyperbolic asymptotically flat spacetimes satisfying the null energy condition [1,3]. Therefore, unless our Universe contains matter violating this energy condition or one is willing to venture into a black hole, any experiment designed to actively detect the Universe's nontrivial topology by sending and receiving causal signals will fail (in the sense that all such signals will be homotopic).

They have also shown that passive topological censorship cannot be expected to hold for all "reasonable" spacetimes by giving an example of a globally hyperbolic asymptotically flat vacuum spacetime having the spatial topology of $\mathbb{R}P^3$ minus a point that violates this condition. Further, combining a theorem due to Schoen and Yau [5] and the results of Gromov and Lawson [6], they argue that any nontrivial topology of a Cauchy surface due to a $K(\pi, 1)$ prime factor [7] must be surrounded by an apparent horizon $\mathcal{A}$, from which they conclude that any violation of passive topological censorship cannot be due to the $K(\pi, 1)$ prime factors.

Were the apparent horizon $\mathcal{A}$ surrounding a $K(\pi, 1)$ factor always a *future* apparent horizon (the expansion of a family of future-directed outgoing null geodesics normal to $\mathcal{A}$ being zero on $\mathcal{A}$), then their argument would be correct as such horizons necessarily lie behind event horizons [4,8]. However, the Schoen-Yau theorem guarantees only that $\mathcal{A}$ will be *either* a future or past apparent horizon, which does not allow one to conclude that these factors in the topology are passively censored.

We show that no such argument can be made by constructing a globally hyperbolic asymptotically flat spacetime satisfying the dominant-energy condition [3], whose Cauchy surfaces are diffeomorphic to the three-torus less a point (T^3 being a $K(\pi, 1)$ manifold), and whose universal covering spacetime violates Eq. (1.2). While we do not show that all $K(\pi, 1)$ prime factors need not be passively censored, there seems to be no reason why they

should. Therefore, while “reasonable” spacetimes must satisfy active topological censorship, a non-simply connected spacetime should not be expected to satisfy passive topological censorship.

Our conventions are those of Ref. [4]. In particular, our spacetime metrics are such that timelike vectors have negative norms.

II. THE COUNTEREXAMPLE

Our counterexample is constructed from a dust-filled $k = 0$ Robertson-Walker spacetime and a Schwarzschild spacetime as follows [9]. (Similar counterexamples whose Cauchy surface topology is a closed hyperbolic space less a point can be constructed from the $k = -1$ Robertson-Walker spacetimes.)

The $k = 0$ Robertson-Walker spacetimes have $\mathbb{R}^3$ Cauchy surfaces and can be coordinated so that the metric h_{ab} is given by

$$h = a^2(\eta)(-d\eta^2 + dx^2 + dy^2 + dz^2) \quad (2.1a)$$

$$= a^2(\eta)(-d\eta^2 + d\chi^2 + \chi^2\Omega), \quad (2.1b)$$

where x , y , and z can be thought of as rectangular coordinates, χ can be thought of as a radial coordinate, and Ω_{ab} is a unit-metric on the two-sphere. With dust as a source, $a(\eta) = C(\eta/2)^2$, for some constant C , and the dust flow is orthogonal to surfaces of constant η . We shall take $\eta > 0$. (Note that these spacetimes are forever expanding to the future.)

From such a $k = 0$ Robertson-Walker spacetime $(\mathbb{R}^4, h_{ab})$, we construct a spacetime with T^3 Cauchy surfaces $(\mathbb{R} \times T^3, h'_{ab})$ by identifying $(\mathbb{R}^4, h_{ab})$ under a discrete group of isometries which, for definiteness, we shall take to be the three translations along x , y , and z by a unit amount. (This way our spacetime is a “unit square cell”.) This spacetime is locally spherically symmetric so it can be locally coordinated by η , χ , and two angular coordinates as in Eq. (2.1b).

We now construct our spacetime (M, g_{ab}) as follows. In the spacetime $(\mathbb{R} \times T^3, h'_{ab})$, consider the (locally spherically symmetric) timelike three-surface $\mathcal{T}$ given by $\chi = \chi_0$ where χ_0 is a constant satisfying $0 < \chi_0 < 1/2$ that will be chosen later. (The upper bound is needed so that this tube “lies within a single cell”.) We now take the region lying *outside* of $\mathcal{T}$ and attach it to a Schwarzschild spacetime of mass $(C/2)\chi_0^3$ by constructing a “similar” timelike three surface $\mathcal{T}'$ therein. See Fig. 1. (The choice of the mass of the Schwarzschild spacetime is found by viewing the Robertson-Walker spacetime as a spherically symmetric spacetime, constructing the mass function m thereon [10], and then evaluating m on $\mathcal{T}$. Further, we claim that such a matching can always be done in such a way that there are no distributional surface layers of matter, i.e., the matching is C^1 .)




FIG. 1. A spacetime diagram representing a maximally extended Schwarzschild spacetime. The curve extending from region III into region IV represents the timelike three-surface $\mathcal{T}'$. The surface $\mathcal{T}'$ is to be attached to its analogous surface $\mathcal{T}$ in the dust-filled spacetime with T^3 Cauchy surfaces. The final spacetime includes everything “outside” of $\mathcal{T}$ in the T^3 Robertson-Walker spacetime and everything “to the right” of $\mathcal{T}'$ in the Schwarzschild spacetime as drawn here.

Our spacetime (M, g_{ab}) is globally hyperbolic having Cauchy surfaces diffeomorphic to T^3 minus a point. Further, the asymptotic region of this spacetime is the asymptotic region I of the Schwarzschild spacetime in Fig. 1. Although this spacetime is not vacuum everywhere, it does satisfy the dominant-energy condition, and indeed just about any reasonable energy condition as the dust has positive energy density and zero pressure. We now show that for an appropriate choice of χ_0 , the spacetime (M, g) does not satisfy passive topological censorship. To do this, we construct its universal covering spacetime and then construct the sets $J^-(\mathcal{J}_i^+)$.

The universal covering spacetime $(\tilde{M}, \tilde{g}_{ab})$ of (M, g_{ab}) can be represented by a $k = 0$ Robertson-Walker spacetime with a Schwarzschild spacetime attached to each timelike three-surface of “radius” χ_0 with “center” having integer x , y , z coordinates. See Fig. 2.




FIG. 2. A 2+1 dimensional version of a portion of the universal covering spacetime of (M, g_{ab}) . Each of the (locally spherically symmetric) timelike three-surfaces are attached to identical (though distinct) Schwarzschild spacetimes. A point p that lies in the overlap region of two downward “cones” can reach distinct null infinities by a future-directed causal curve. Therefore, there exist homotopically distinct future-directed causal curves from p to $\mathcal{J}^+$.

Consider the null infinity $\mathcal{J}_i$ associated with one of the asymptotic Schwarzschild regions, e.g., the one “centered” at $(x, y, z) = (0, 0, 0)$. From Fig. 1, we see that the portion of $J^-(\mathcal{J}_i^+)$ in the Robertson-Walker part of the spacetime is equivalent to the past of the portion of $\mathcal{T}'_i$ which lies in region III of the Schwarzschild spacetime in Fig. 1. This is the portion of $\mathcal{T}'_i$ (equivalently $\mathcal{T}_i$) where $\nabla^a r$ is past-directed timelike, where r is the size

of the spheres of symmetry, which in our case is given by $r = a(\eta)\chi$. A simple calculation shows that $\nabla^a r$ is past-directed timelike on the region where $\eta < 2\chi$. Therefore, the region we are interested in is that shaded in Fig. 3. For a point p in this region, there exists a future-directed causal curve passing through $\mathcal{T}_i$ that connects p to $\mathcal{J}_i^+$.




FIG. 3. A spacetime diagram showing the portion of $J^-(\mathcal{J}_i^+)$ in the Robertson-Walker part of the spacetime (i.e., the region shaded). The vertical line at $\chi = \chi_0$ represents $\mathcal{T}_i$. The region of interest is precisely the region that is both “outside” of $\mathcal{T}_i$ and that lies to the past of the portion of $\mathcal{T}_i$ on which $\nabla^a r$ is past-directed timelike. (In this diagram, each point represents a sphere, excepting along the line $\chi = 0$.)

So, since our “Schwarzschild ends” in the universal covering spacetime are “separated by a unit amount”, from Fig. 3 it is clear that if we construct our spacetime with $\chi_0 > 1/6$, the “shaded regions” will overlap, and therefore, as illustrated in Fig. 2, the pasts of distinct future null infinities will overlap. Therefore, condition Eq. (1.2) is violated for $\chi_0 > 1/6$, showing that these spacetimes violate passive topological censorship.

ACKNOWLEDGMENTS

I would like to thank John Friedman and Donald Witt for their comments and for answering my many questions.

- [5] R. Schoen and S. -T. Yau, *Commun. Math. Phys.* **79**, 231 (1981).
- [6] M. Gromov and H. B. Lawson, Jr., *Inst. Hautes Etudes Sci. Publ. Math.* **58**, 83 (1983).
- [7] For a discussion of the prime factorizations of three-manifolds, see J. Hempel, *3-Manifolds* (Princeton University Press, Princeton, 1976). A closed manifold Σ is a $K(\pi, 1)$ manifold, for some group π , if $\pi_1(\Sigma) = \pi$ (i.e., the fundamental group of Σ is the group π) and $\pi_k(\Sigma) = 0$ for $k \neq 1$ (i.e., all homotopy groups of Σ higher than the first are trivial). Equivalently, a closed manifold Σ is a $K(\pi, 1)$ manifold if its universal covering manifold is deformable to a point. For example, the three-torus T^3 , having $\mathbb{R}^3$ as its universal cover, is a $K(\pi, 1)$ manifold.
- [8] S. W. Hawking and G. F. R. Ellis, *The Large-Scale Structure of Space-time* (Cambridge University Press, Cambridge, England, 1973).
- [9] D. R. Brill, in *Proceedings of the Third Marcel Grossmann Meeting on the Recent Developments of General Relativity*, edited by H. Ning (North Holland, New York, 1983).
- [10] G. A. Burnett, *Phys. Rev. D* **48**, 5688 (1993).

-
- [1] J. L. Friedman, K. Schleich, and D. M. Witt, *Phys. Rev. Lett.* **71**, 1486 (1993).
 - [2] J. L. Friedman, in *Conceptual Problems of Quantum Gravity*, edited by A. Ashtekar and J. Stachel (Birkhauser, Boston, 1991).
 - [3] A spacetime with Einstein tensor G_{ab} is said to satisfy the null energy condition (also known as the null-convergence condition) if $G_{ab}k^a k^b \geq 0$ for all null k^a . Actually, they prove their theorem under a slightly weaker condition known as the averaged null energy condition. The dominant-energy condition is said to be satisfied if $G_{ab}t^a u^b \geq 0$ for all future-directed timelike t^a and u^b .
 - [4] R. M. Wald, *General Relativity* (University of Chicago Press, Chicago, 1984).